

What is claimed is:

1. An isolated polypeptide comprising:
  - a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-2,
  - 5        b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-2,
  - c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-2, or
  - d) an immunogenic fragment of an amino acid sequence selected from the group consisting  
10      of SEQ ID NO:1-2.
2. An isolated polypeptide of claim 1, having an amino acid sequence selected from the group consisting of SEQ ID NO:1-2.
- 15        3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide of claim 3, having a sequence selected from the group consisting of SEQ ID NO:3-4.
- 20        5. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
6. A cell transformed with a recombinant polynucleotide of claim 5.
- 25        7. A transgenic organism comprising a polynucleotide of claim 5.
8. A method for producing a polypeptide of claim 1, the method comprising:
  - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide  
30      comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
  - b) recovering the polypeptide so expressed.
9. An isolated antibody which specifically binds to a polypeptide of claim 1.

10. An isolated polynucleotide comprising:
- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:3-4,
  - b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:3-4,
  - 5 c) a polynucleotide sequence complementary to a), or
  - d) a polynucleotide sequence complementary to b).

11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.

12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:

- a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.

14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.

15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

16. A method of treating a disease or condition associated with decreased expression of functional CIRYP, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.

17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

19. A method of treating a disease or condition associated with decreased expression of functional CIRYP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

b) detecting antagonist activity in the sample.

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with overexpression of functional CIRYP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

a) exposing a sample comprising the target polynucleotide to a compound, and

b) detecting altered expression of the target polynucleotide.